

Which is the solutions quality of a LS ?


- for a given neighborhood
- even in exponential time !

Grover ‘‘92’’

- TSP with 2 – exchange
- Bipartitioning of a weighted graph with 2 – exchange

$$C_{\text{loc}} \leq C_{AV}$$

local minimum average cost of all solutions



The Quadratic Assignment Problem (QAP)

- n locations : distance
- n facilities: flow f_{ij}
- $\pi(i)=k$: facility i \rightarrow location k

minimize the total cost


$$\text{Min}_{\pi \in \Pi} \sum_{i,k=1}^n f_{ik} d_{\pi(i)\pi(k)}$$


The QAP: an example

$\pi = 3 \ 1 \ 2 \ 5 \ 4 \ 6$

$\pi_{ij} = 3 \ 4 \ 2 \ 5 \ 1 \ 6$

2-exch.





The QAP


→ PLS-complete(neighb. 2-exchange)
(schaffer & Yannakakis '91)

2-exchange:

$$\pi = (\pi(1), \dots, \pi(i), \dots, \pi(j), \dots, \pi(n))$$

$$\pi_{ij} = (\pi(1), \dots, \pi(j), \dots, \pi(i), \dots, \pi(n))$$

$$N(\pi) = (n * (n - 1)) / 2$$



Notations

- $S(F) = \sum_{i,j} f_{ij}$
- $x = (x_1, x_2, \dots, x_n)$
 $y = (y_1, y_2, \dots, y_n)$
 $\pi y = (y_{\pi(1)}, y_{\pi(2)}, \dots, y_{\pi(n)})$
 $\langle x, y \rangle_+ = \max_\pi \langle x, \pi y \rangle$
 $\langle x, y \rangle_- = \min_\pi \langle x, \pi y \rangle$
- $F_k = \sum_i f_{ki} = \sum_i f_{ik}$
- $C_{AV} = \frac{1}{n!} \sum C(\Pi)$
- $C_{\overline{123}}$

Solution quality for the QAP ?

E.Angel & V.Z.'96 '' (DAM 98)

Th₁ :

$$C_{\overline{\text{loc}}} \leq \frac{\langle F, D \rangle_+}{S(F)S(D)} n C_{\text{AV}} = \frac{\langle F, D \rangle_+}{2(n-1)} \leq \frac{n}{2} C_{\text{AV}}$$

Th₂ :

$$\leq \frac{n}{2} C_{\text{AV}} \text{ in } O\left(n \log \frac{S(F) \times S(D)}{2}\right) \text{ iterat.}$$

Idea (deepest local search)

$$\delta_{ij}(\pi) = C(\pi_{ij}) - C(\pi)$$

$$\nabla^2 C(\pi) = \frac{1}{n(n-1)} \sum_{i,j} \delta_{ij}(\pi) \leq -\frac{4}{n} (C(\pi) - C_{AV}) + a$$

$\forall \pi \exists i, j \text{ t. q.}$

$$\delta_{ij}(\pi) \leq \nabla^2 C(\pi)$$

$$C(\pi) \rightarrow C(\pi') \leq C(\pi) - \frac{4}{n} (C(\pi) - C_{AV}) + a$$

$$1 = 1 - \frac{4}{n} (1 - C_{AV}) + a$$

$$C_{AV} = \frac{S(F) \times S(D)}{2n(n-1)}$$

$$1) \frac{C_{\max}}{\frac{n}{2} C_{AV}} \rightarrow \infty$$

e_x.

$$F = D = \begin{pmatrix} 0 & 1 & 1 & \cdot & \cdot & \cdot & 1 \\ 1 & 0 & 1 & \cdot & \cdot & \cdot & 1 \\ \cdot & 1 & 0 & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & 0 & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & n^2 \\ 1 & \cdot & \cdot & \cdot & 1 & n^2 & 0 \end{pmatrix}$$

$$\frac{n}{2} C_{AV} = \frac{(n - 1)(3n + 2)^2}{4}$$

$$C_{\max} = \frac{n(n - 1)}{2} - 1 + n^4$$

2) Traveling Salesman Problem (2-exchange)

- $D = (d_{ij})$ adj. matrix

$$\bullet F = (f_{ij}) = \begin{pmatrix} 0 & 1 & 0 & . & . & . & . & . & 0 & 1 \\ 1 & 0 & 1 & 0 & . & . & . & . & . & 0 \\ 0 & 1 & 0 & 1 & 0 & . & . & . & . & . \\ 0 & 0 & 1 & 0 & 1 & 0 & . & . & . & . \\ . & . & 0 & 1 & 0 & 1 & 0 & . & . & . \\ . & . & . & 0 & 1 & 0 & 1 & 0 & . & . \\ . & . & . & . & 0 & 1 & 0 & 1 & 0 & 0 \\ . & . & . & . & . & 0 & 1 & 0 & 1 & 0 \\ 0 & . & . & . & . & . & 0 & 1 & 0 & 1 \\ 1 & 0 & . & . & . & . & . & 0 & 1 & 0 \end{pmatrix}$$

$$S(F) = 2n$$

$$\langle F, D \rangle_+ = 2S(D)$$


Particular cases (TSP)

Th₁ ⇒ conclusion :

$$C_{\overline{\text{loc}}} \leq C_{AV}$$

[Grover]

$$C_{\overline{\text{loc}}} \leq \frac{\langle F, D \rangle_+}{S(F)S(D)} n C_{AV}$$



3) Bipartitioning

- weighted graph $G(V,E)$

- 2-exchange

$$|A| = |B| = \frac{|V|}{2} (\pm 1)$$

– $D = (d_{ij})$ adj. matrix

Particular cases (Bipartitioning)

$$F = \begin{pmatrix} f_{ij} \end{pmatrix} = \begin{pmatrix} O & U \\ U & O \end{pmatrix} \xrightarrow{\leftrightarrow} n/2$$

$$U = \begin{pmatrix} u_{ij} \end{pmatrix} = \begin{cases} 1, & i, j = 1, \dots, \frac{n}{2} \\ 0 & \text{otherwise} \end{cases}$$


$$\boxed{C_{\text{loc}} \leq C_{\text{AV}}}$$

$$C_{\overline{\text{loc}}} \leq \frac{\langle F, D \rangle_+}{S(F)S(D)} n C_{\text{AV}}$$

[Grover]

$\frac{n}{2} S(D)$ ↗

$n \frac{n}{2}$ ↗



m-densest (m-lightest)

- Graph $G(V, E)$, $|V| = n$

consider $m (\leq n)$, $m = 1, \dots, n$

"Find a sub - graph $G'_G(V', E')$ with $|V'| = m$

and $|E'|$ maximum (minimum)

- NP - complete

- maximum clique
- maximum Independent set

Results (2-exchange)

$$F = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{matrix} \updownarrow \\ m \\ \updownarrow \\ n-m \end{matrix} D = \begin{pmatrix} d_{ij} \end{pmatrix}$$

$d_1 \geq d_2 \geq \dots \geq d_n$ degrees of G

$$C_{\text{loc}}^+ \geq \frac{(m - 1)}{2(n - 1)} (d_n + d_{n-1} + \dots + d_{n-m+1})$$

$$C_{\text{loc}}^- \leq \frac{(m - 1)}{2(n - 1)} (d_1 + d_2 + \dots + d_m)$$

$$\frac{T_h}{\overline{\overline{h}}_1} \left\{ \begin{array}{l} S(F) = m(m - 1) \\ S(D) = 2|E| \\ C_{\text{av}} = \frac{m(m - 1)}{n(n - 1)} |E| \\ \langle F, D \rangle_+ = (m - 1)(d_1 + \dots + d_m) \end{array} \right.$$

5) Stable Maximum

(Meyers " 72 ")

" If $d_1 \geq d_2 \geq \dots \geq d_n$ and $d_1 + d_2 + \dots + d_k \leq n - k$,
for $2 \leq k \leq n$ then stable maximal $\geq k$ "

(Angel & V.Z. " 96 ")

" If $d_1 + d_2 + \dots + d_k \leq \left\lfloor \frac{2a(n-1)}{k-1} \right\rfloor_*$ for a integer,

$2 \leq k \leq n$

$$\lfloor x \rfloor_* = \begin{cases} x - 1 & \text{if } x \text{ integer} \\ \lfloor x \rfloor & \text{otherwise} \end{cases}$$

stable maximal $\geq k - a + 1$ "

Example

$n=10$


3 3 3 2 2 2 1 1 1


$6 \leq 10 - 2$ Meyer

$a=3$

$K=5$

$k-a+1=3$






Comparison

- Meyer's result is deductive
 - This result is constructive
 - This condition is more flexible
 - Result interesting in practice
- " C_{loc} includes at most $a - 1$ edges "
- we remove at most $a - 1$ vertices
- for obtaining a maximal Independent set


Example


ex

- $a=5$
4 edges
4 vertices




- $a=7$
6 edges
3 vertices






LS Drawbacks

- Local optimum
- “good“ neighborhoods
- exploration strategies
- Performances guarantee ?
- Parallelization ?



Conclusions

- Some new results on the performance of Local Search and its variants
- The k-densest and the k-lightest problems
- A constructive result for the max. Ind. Set Pr.
(Th. MEYER)



FUTURE WORK

- ?!
 - Proposed bound for QAP is it attained?
 - TSP with 2-OPT is it in PLS - complete?
 - Other problems with guaranteed LS
 - Parallelization of PBs in PLS
 - `` Good '' neighborhood structures
 - Scheduling scheme in SA
- }